arXiv:1603.07874v1 [math.RT] 25 Mar 2016

ON SHALIKA GERMS

DAVID KAZHDAN

Abstract. Let G be a reductive group over a local field F satisfying the assumptions of [Deb1], $G_{reg} \subset G$ the subset of regular elements. Let $T \subset G$ be a maximal torus. We write $T_{reg} = T \cap G_{reg}$. Let dg, dt be Haar measures on G and T. They define an invariant measure dg/dt on G/T. Let $\mathcal{H}$ be the space of complex valued locally constant functions on G with compact support. For any $f \in \mathcal{H}, t \in T_{reg}$ we define $I_t(f) = \int_{G/T} f(\bar{g}t\bar{g}^{-1})dg/dt$. Let P be the set of conjugacy classes of unipotent elements in G. For any $\Omega \in P$ we fix an invariant measure ω on Ω. As well known [R] for any $f \in \mathcal{H}$ the integral

$$I_\Omega(f) = \int_\Omega f\omega$$

is absolutely convergent. Shalika [Sh] has shown that there exist functions $\tilde{j}_\Omega(t), \Omega \in P$ on $T \cap G_{reg}$ such that

$$I_t(f) = \sum_{\Omega \in P} \tilde{j}_\Omega(t) I_\Omega(f) \qquad (\star)$$

for any $f \in \mathcal{H}, t \in T$ *near* to e where the notion of *near* depends on f.

For any positive real number r one defines an open Ad-invariant subset G_r of G and a subspace $\mathcal{H}_r$ as in [Deb1]. In this paper I show that for any $f \in \mathcal{H}_r$ the equality $(\star)$ is true for all $t \in T_{reg} \cap G_r$.

Let F be a local non-archimedian field, $\mathcal{O}$ be the ring of integers of $F, \mathfrak{m} \subset \mathcal{O}$ the maximal ideal, $k = \mathcal{O}/\mathfrak{m}$ the residue field, ζ a generator of $\mathfrak{m}, q = |k|$ and $val : F^\star \to \mathbb{Z}$ the valuation such that $val(\zeta) = 1$.

Let G a reductive F-group, $G_{reg} \subset G$ the subset of regular elements. To simplify notations I write G instead of $G(F)$ and fix a Haar measure dg on G. For any $r \in \mathbb{Z}_+$ we denote by G_r the Ad-invariant open subset as in [Deb1].

Let $\mathfrak{g}$ be the Lie algebra of G and $\mathfrak{g}_{nil} \subset \mathfrak{g}$ be the subset of elements x of the form $x = n + z$ where n is a topologically nilpotent element of $[\mathfrak{g}, \mathfrak{g}]$ and $z \in Z_{\mathfrak{g}}(\mathfrak{m})$ where $Z_{\mathfrak{g}}$ is the center of $\mathfrak{g}$.

Let $\mathcal{H} = \mathcal{S}(G)$ be the space of locally constant $\mathbb{C}$-valued compactly supported functions on G. Since we fixed a Haar measure on $G, \mathcal{H}$ has a natural algebra structure. We denote by $\bar{\mathcal{H}}$ the space of locally constant $\mathbb{C}$-valued compactly supported function on $\mathfrak{g}$.

I will freely use notations and results of [Deb1]. In particular we fix an map $\phi : \mathfrak{g}_n \to G$ as in [Deb1]. (So $\phi(x) = Id_n + x$ if $G = Gl_n$ and ϕ is the inverse of the quasi-logarithm map (see [BKV, Appendix C]) if

Date: March 28, 2016.

$char(k) >> 1$.) and consider open Ad-invariant subsets $G_r \subset G, \mathfrak{g}_r \subset \mathfrak{g}$ as in [Deb1]. Then $\phi(\mathfrak{g}_r) = G_r$ (see [BKV, Lemma C.4]).

We fix a maximal torus T in G and a Haar measure dt on T. Let $\mathfrak{t}$ be the Lie algebra of $T, T_{reg} \subset T, \mathfrak{t}_{reg} \subset \mathfrak{t}$ the subsets of regular elements.

For any regular $t \in T_{reg}, f \in \mathcal{H}$ we define

$$I_t(f) = \int_{G/T} f(g\phi(t)g^{-1})dg/dt.$$

Analogously for any regular $\bar{t} \in \mathfrak{t}_{reg}, \bar{f} \in \bar{\mathcal{H}}$ we define

$$I_{\bar{t}}(\bar{f}) = \int_{G/T} \bar{f}(g\bar{t}g^{-1})dg/dt.$$

It is clear that for any $f \in \mathcal{H}, \bar{t} \in \mathfrak{t}_{reg} \cap \mathfrak{g}_{nil}$ we have

$$I_t(f) = I_{\bar{t}}(\phi^\star(f)), t = \phi(\bar{t}).$$

We define $\mathcal{H}_r \subset \mathcal{H}$ as the subspace of $f \in \mathcal{H}$ such that $\pi(f) = 0$ for all irreducible representations π of G of $depth > r$. As follows from [Deb2, Teorem 5.5.4] this subspace coincides with the subspace $\mathcal{H}_r$ of [Deb1].

Let P be the set of conjugacy classes of nilpotent elements in G. For any $\Omega \in P$ we fix an invariant measure ω on Ω. As well known [R] for any $\bar{f} \in \bar{\mathcal{H}}$ the integral

$$I_\Omega(\bar{f}) = \int_\Omega \bar{f}\omega$$

is absolutely convergent and $I_\Omega(\bar{f}_\zeta) = q^d I_\Omega(\bar{f})$ where $\bar{f}_\zeta(x) = f(\zeta^2 x)$.

The following result is proven in [Deb1, Conjecture 2].

Claim. For any $f \in \mathcal{H}_r$ such that $I_\Omega(\phi^\star(f)) = 0$ for all $\Omega \in P$ we have $I_t(f) = 0$ for all $t \in T_{reg} \cap G_r$.

Shalika [Sh] has shown that there exist functions $\tilde{j}_\Omega(t), \Omega \in P$ on $T \cap G_{reg}$ such that

$$I_t(f) = \sum_{\Omega \in P} \tilde{j}_\Omega(t) I_\Omega(f)$$

for any $t \in T$ near to e. Let j_Ω be the function on a neighboorhood of 0 in $\mathcal{T}$ given by $j_\Omega(x) = \tilde{j}_\Omega(\phi(x))$.

Harish-Chandra [HC] has shown that

$$j_\Omega(c^2 x) = j_\Omega(x)|c|^{dim(\mathcal{N}) - dim(\Omega)}$$

for all $c \in F^\star$ if $x \in \mathcal{T}$ is near to 0.

Using this homogenuity we can define functions $j_\Omega(t)$ for all $t \in \mathcal{T}$.

Theorem. For any $f \in \mathcal{H}_r$ the Shalika germ expansion is true for all $g \in \mathcal{T} \cap \mathfrak{g}_r$.

Proof. For any $\Omega \in P$ we define $\mathcal{H}_r(\Omega)$ to be the set of all $f \in \mathcal{H}_r$ such that $I_{\Omega'}(f) = 0$ for all $\Omega' \in P$ different from Ω.

As follows from [Deb1, Theorem 4.1.4] the space $\mathcal{H}_r$ is spanned by subspaces $\mathcal{H}_r(\Omega), \Omega \in P$. So it is sufficient for a proof of Theorem to show that

$$q^d I_{\bar{t}}(\bar{f}) = I_{\zeta^2\bar{t}}(\bar{f}), \bar{f} = \phi^\star(f)$$

for all $f \in \mathcal{H}_r(\Omega)$ all regular $\bar{t} \in \mathfrak{t} \cap \mathfrak{g}_r$.

Let

$$\bar{f} = \phi^\star(f), h = q^d \bar{f} - \bar{f}_\zeta$$

where $d = dim(\Omega)$. Then $I_{\Omega'}(h) = 0$ for all $\Omega' \in P$ and it follows from the Claim that $I_t(h) = 0$ for all $t \in \mathcal{T}_r$. So

$$q^d I_{\bar{t}}(\bar{f}) = I_{\zeta^2\bar{t}}(\bar{f}), \bar{f} = \phi^\star(f).$$

References

[BKV] Roman Bezrukavnikov, David Kazhdan and Yakov Varshavsky, *On the depth r Bernstein projector*, arXiv:1504.01353.

[Deb1] Debacker, Stephen, *Homogeneity results for invariant distributions of a reductive p-adic group*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 3, 391–422.

[Deb2] Debacker, Stephen, *Some applications of Bruhat-Tits theory to harmonic analysis on a reductive p-adic group*, Michigan Math. J. **50**, Issue 2 (2002), 241–261.

[HC] Haris-Chandra, *Admissible invariant distributions on reductive p-adic groups*, Lie theories and their applications (Proc. Ann. Sem. Canad. Math. Congr., Queen's Univ., Kingston, Ont., 1977), pp. 281–347. Queen's Papers in Pure Appl. Math., No. 48.

[R] Rao, R., *Orbital integrals in reductive groups*, Annales of Math. (2) **96** (1972), 505–510.

[Sh] Shalika, J. A., *A theorem on semi-simple P-adic groups*, Ann. of Math. (2) **95** (1972), 226–242.

Institute of Mathematics, The Hebrew University of Jerusalem, Givat-Ram, Jerusalem, 91904, Israel

E-mail address: kazhdan@math.huji.ac.il